

WE CLAIM:

1. A method for hydrolyzing α -galactose bonds comprising:
contacting a compound having an α -galactose bond with an effective amount of
an enzyme having at least a 70% amino acid identity to amino acid sequence set forth in SEQ ID

NO: 4.

2. The method according to claim 1 wherein the enzyme has at least 90% amino acid
identity to the amino acid sequence set forth in SEQ ID NO: 4.

3. The method according to claim 1 wherein the enzyme comprises a sequence of
amino acids identical to amino acids 1 to 364 of SEQ ID NO:4.

4. The method according to claim 1 wherein the enzyme has the amino acid
sequence as set forth in SEQ ID NO: 4.

5. The method according to claim 1 wherein the enzyme is recombinantly produced.

6. The method according to claim 1 wherein the compound having the α -galactose
bond is raffinose.

7. The method according to claim 6 wherein the α -galactose bond is in raw beet
sugar.

8. The method according to claim 1 wherein the compound is raffinose, stachyose,
verbascose, or a combination thereof.

9. The method according to claim 8 wherein the compound is contained in a member
of the lentil or bean family, or a combination thereof.

10. The method according to claim 1 wherein the contacting is at a temperature of about 85° C.

11. The method according to claim 1 wherein the contacting is at a pH of about 9.5.

12. The method according to claim 1 wherein the contacting is at a temperature of about 85° C and a pH of about 9.5.

Add c5

006T40" 2806T960